

Markov Properties of the Fluctuation Limit of a Particle System with Death*

BEGOÑA FERNÁNDEZ

*Facultad de Ciencias, UNAM, México 04510, D. F. México and
Centro de Investigación y de Estudios Avanzados, México 07000, D. F. México*

Submitted by David Aldous

Received May 8, 1989

Consider a sequence of independent Brownian motions in R^d whose initial positions are distributed according to a probability measure μ on $R^d \times R_+$ and disappear after an exponentially distributed lifetime with parameter λ . It is shown that the fluctuation process of the empirical distribution of the first n motions converges weakly in the Skorokhod spaces $D([0, T_0], \mathcal{S}'(R^d))$ and $D([T_0, \infty), \mathcal{S}'(R^d))$ as $n \rightarrow \infty$, where $T_0 = \inf\{t : \mu(R^d \times [t, \infty)) = 0\}$, to a continuous, centered Gaussian process X . Conditions for X to be Markovian are determined in terms of μ . The cases $\lambda > 0$ and $\lambda = 0$ (i.e., the motions go on forever) exhibit different Markovian properties. © 1991 Academic Press, Inc.

1. INTRODUCTION

Let $B^k = \{B_t^k, t \geq 0\}$, $k = 1, 2, \dots$, be independent Brownian motions in R^d , each one starting from a point distributed according to a probability measure μ on $R^d \times R_+$, and disappearing after an exponentially distributed lifetime with parameter λ . In this paper we study the asymptotic behavior of the fluctuation process of the empirical distribution of the first n motions as $n \rightarrow \infty$, and the Markov property of the fluctuation limit process. The case $\lambda = 0$ (i.e., the motions go on forever) was considered in [6], and the case $\lambda = 0$ and $\mu(R^d \times \{0\}) = 1$ was studied by Itô [7]. In contrast with the case $\lambda = 0$, where the fluctuation limit process is Markovian, in the model with death the Markov property does not hold in general.

The processes we consider can be regarded as $\mathcal{S}'(R^d)$ -valued, where $\mathcal{S}'(R^d)$ is the Schwartz space of tempered distributions, i.e., the topological dual of the space $\mathcal{S}(R^d)$ of rapidly decreasing infinitely differentiable real functions on R^d . The duality between these two spaces will be denoted by $\langle \cdot, \cdot \rangle$.

* Research partially supported by CONACyT Grant PCEXCNA-040319.

We will prove that the fluctuation processes of the empirical distribution of the first n motions converge weakly in the Skorokhod spaces $D([0, T_0], \mathcal{S}'(R^d))$ and $D([T_0, \infty), \mathcal{S}'(R^d))$ as $n \rightarrow \infty$, where $T_0 = \inf\{t: \mu(R^d \times [t, \infty)) = 0\}$, to a continuous, centered Gaussian process X , and that X has the Markov property in an interval $[a, b]$ if and only if $\lambda = 0$ or $\mu(R^d \times (a, b]) = 0$. Thus the Markov property does not hold for X in $[a, b]$ if and only if births and deaths are both possible in $(a, b]$. This result should be considered natural since it is in agreement with the fact that the same Markovian behavior holds for a single Brownian motion with initial distribution μ and exponential lifetime. We will also prove that if $\mu(R^d \times (a, b]) = 0$ then the process X has the *strict Markov property* (defined below) on $(a, b]$, and we will obtain the Langevin equation for X on $(a, b]$. In particular the process $\{X_{T_0+t}, t \geq 0\}$ is strictly Markovian.

We will now recall the Markov property and the *strict Markov Property* for Gaussian $\mathcal{S}'(R^d)$ -valued processes (see [6]).

The markov property of an $\mathcal{S}'(R^d)$ -valued continuous, centered, Gaussian process X is equivalent to the statement that given $s < t$ there exists an $\mathcal{S}'(R^d)$ -valued random variable $X_{s,t}$ such that for each $\psi \in \mathcal{S}(R^d)$, $\langle X_{s,t}, \psi \rangle$ belongs to $\langle X_s, \mathcal{S}(R^d) \rangle^-$ (the L^2 -closure of the linear space of random variables $\{\langle X_s, \phi \rangle, \phi \in \mathcal{S}(R^d)\}$), and

$$\text{Cov}(\langle X_r, \phi \rangle, \langle X_t, \psi \rangle) = \text{Cov}(\langle X_r, \phi \rangle, \langle X_{s,t}, \psi \rangle) \quad (\text{M})$$

for all $r \leq s$ and $\psi, \phi \in \mathcal{S}(R^d)$. We define the *strict Markov property* of a centered, Gaussian $\mathcal{S}'(R^d)$ -valued process as follows: given $s < t$ there exists a continuous linear mapping $\mathcal{U}_{s,t}: \mathcal{S}(R^d) \rightarrow \mathcal{S}(R^d)$ such that

$$\text{Cov}(\langle X_r, \phi \rangle, \langle X_t, \psi \rangle) = \text{Cov}(\langle X_r, \phi \rangle, \langle X_s, \mathcal{U}_{s,t}\psi \rangle) \quad (\text{SM})$$

for all $r \leq s$ and $\psi, \phi \in \mathcal{S}(R^d)$. Clearly this is a special case of the Markov property since $\langle X_s, \mathcal{U}_{s,t}\psi \rangle$ belongs to $\langle X_s, \mathcal{S}(R^d) \rangle^-$. If $\{\mathcal{U}_{s,t}, s \leq t\}$ is a reverse evolution system on $\mathcal{S}(R^d)$, i.e., $\mathcal{U}_{t,t} = I$ and $\mathcal{U}_{r,s}\mathcal{U}_{s,t} = \mathcal{U}_{r,t}$ for $r \leq s \leq t$, then it is easy to see that (SM) is equivalent to

$$\text{Cov}(\langle X_s, \phi \rangle, \langle X_t, \psi \rangle) = \text{Cov}(\langle X_s, \phi \rangle, \langle X_s, \mathcal{U}_{s,t}\psi \rangle). \quad (\text{SM}')$$

The topologies on $\mathcal{S}(R^d)$ and $\mathcal{S}'(R^d)$ are well known (see [6]), and we will denote by $\mathcal{S}(R^d) \oplus R$ the direct sum of $\mathcal{S}(R^d)$ and the space of constant functions on R^d with the direct sum topology.

The space of right-continuous with left limits functions from $[a, b]$ into $\mathcal{S}'(R^d)$ is denoted by $D([a, b], \mathcal{S}'(R^d))$ and is endowed with a Skorokhod-type topology [9].

The following notation will be used throughout:

$$p_t(x, y) = (2\pi t)^{-d/2} \exp\{-|y - x|^2/2t\}, \quad t > 0, x, y \in R^d.$$

$$\mathcal{T}_t \phi(x) = \int_{R^d} \phi(y) p_t(x, y) dy, \quad t > 0, \mathcal{T}_0 = I \text{ (the Brownian semigroup)}.$$

For $s \leq t$,

$$\begin{aligned} \mu_{st}(x) &= \int_{R^d \times [0, s] \times [t, \infty]} p_{t-u}(x, y) v(dy, du, dl), \\ f_{st}(\phi) &= \int_{R^d \times [0, s] \times [t, \infty]} \phi(x) v(dx, du, dl), \end{aligned}$$

where

$$v(dx, dr, dl) = \begin{cases} \lambda \exp\{-\lambda(l-r)\} dl \mu(dx, dr), & l \geq r, \\ 0, & l < r, \end{cases}$$

if $\lambda > 0$ and

$$v(dx, dr, dl) = \delta_{+\infty}(dl) \mu(dx, dr)$$

if $\lambda = 0$, where $\delta_{+\infty}$ is the Dirac measure concentrated at $+\infty$. Observe that for $\lambda = 0$ the above expressions satisfy

$$\mu_{st}(x) = \int_{R^d \times [0, s]} p_{t-u}(x, y) \mu(dy, du), \quad s \leq t,$$

and

$$f_{st}(\phi) = \int_{R^d \times [0, s]} \phi(x) \mu(dx, du), \quad s \leq t.$$

Then, for $\lambda = 0$ we have $\mu_{st}(x) \equiv \mu_s(x)$ and $f_{st}(\phi) \equiv f_s(\phi)$, where the right-hand sides are the notation used in [6].

The definitions above are valid for bounded, measurable, real functions ϕ on R^d .

Let $T_0 = \inf\{t: \mu(R^d \times [t, \infty)) = 0\}$. We will assume that the measure μ is sufficiently smooth so that for each $T < T_0$,

$$\int_{R^d \times \{s, t\}} e^{-\lambda(t-u)} \mu(dx, du) \leq K(t-s)$$

for all $s < t \leq T$ and some constant K depending on T .

2. RESULTS AND PROOFS

Let $W \equiv \{W_t, t \geq 0\}$ be a standard Brownian motion on R^d (starting from 0), and (Z, τ, θ) and $R^d \times R_+ \times R_+$ -valued random variable with distribution $v(dx, dr, dl)$ independent of W . Let $W^k, (Z^k, \tau^k, \theta^k), k = 1, 2, \dots$, be independent copies of $W, (Z, \tau, \theta)$. We consider as state space $R^d \cup \{\gamma\}$, where γ is an external point, and define the k th Brownian motion with distribution μ and exponential lifetime (i.e., beginning at time τ^k and position Z^k and dying at time θ^k) as

$$B_t^k = \begin{cases} \gamma, & \text{if } t < \tau^k \text{ or } t \geq \theta^k, \\ Z^k + W_{t-\tau^k}^k, & \text{if } \tau^k \leq t < \theta^k. \end{cases}$$

Let N_t^n denote the empirical distribution at time t of the first n processes, i.e.,

$$N_t^n = \sum_{k=1}^n \delta_{B_t^k}, \quad t \geq 0,$$

where δ_x is the Dirac measure concentrated at x . $N^n \equiv \{N_t^n, t \geq 0\}$ can be regarded as an $\mathcal{S}'(R^d)$ -valued process given by

$$\langle N_t^n, \phi \rangle = \sum_{k=1}^n \phi(B_t^k), \quad \phi \in \mathcal{S}(R^d), \quad (2.1)$$

where the definition of ϕ is extended to γ by $\phi(\gamma) = 0$.

We define the fluctuation process $X^n \equiv \{X_t^n, t \geq 0\}$ by centering and normalizing

$$\langle X_t^n, \phi \rangle = n^{-1/2} \sum_{k=1}^n [\phi(B_t^k) - E\phi(B_t^k)], \quad \phi \in \mathcal{S}(R^d). \quad (2.2)$$

Note that (2.1) and (2.2) are also well defined for $\phi \in \mathcal{S}(R^d) \oplus R$.

LEMMA 2.1. For each $t \geq 0$ and $\phi \in \mathcal{S}(R^d) \oplus R$,

$$E\langle N_t^n, \phi \rangle = n \int_{R^d} \phi(y) \mu_n(y) dy, \quad (2.3)$$

and for $s \leq r \leq t$, $\phi, \psi \in \mathcal{S}(R^d) \oplus R$,

$$\begin{aligned}
& \text{Cov}(\langle N_s^n, \phi \rangle, \langle N_t^n, \psi \rangle) \\
&= n \left\{ \int_{R^d} \phi(z) \mu_{ss}(z) e^{-\lambda(t-s)} \mathcal{T}_{t-s} \psi(z) dz \right. \\
&\quad - \int_{R^d} \phi(z) \mu_{ss}(z) dz \left[\int_{R^d} \mu_{ss}(z) e^{-\lambda(t-s)} \mathcal{T}_{t-s} \psi(z) dz \right. \\
&\quad \left. \left. + \int_{R^d \times (s, t]} e^{-\lambda(t-r)} \mathcal{T}_{t-r} \psi(x) \mu(dx, dr) \right] \right\}. \quad (2.4)
\end{aligned}$$

Proof. By (2.1),

$$E\langle N_t^n, \phi \rangle = \sum_{k=1}^n E\phi(B_t^k).$$

Conditioning with respect to (Z^k, τ^k, θ^k) and using Fubini's theorem we have

$$\begin{aligned}
E\phi(B_t^k) &= \int_{R^d \times [0, t] \times [t, \infty)} \mathcal{T}_{t-r} \phi(x) v(dx, dr, dl) \\
&= \int_{R^d} \phi(y) \mu_{tt}(y) dy, \quad k = 1, 2, \dots,
\end{aligned}$$

which proves (2.3).

Using the independence of the B^k , $k \geq 1$, a similar calculation yields (2.4).

We will now state and prove the fluctuation limit theorem.

THEOREM 2.1. *There exists a continuous centered Gaussian $\mathcal{S}'(R^d)$ -valued process $X \equiv \{\langle X_t, \phi \rangle, t \geq 0, \phi \in \mathcal{S}(R^d)\}$ with covariance functional given by the right-hand side of (2.4) with $n = 1$, and X^n converges weakly as $n \rightarrow \infty$ to X in $D([0, T_0], \mathcal{S}'(R^d))$ and $D([T_0, \infty), \mathcal{S}'(R^d))$.*

Proof. The stated process X exists by Kolmogorov's extension theorem for $\mathcal{S}'(R^d)$ -valued processes.

By Theorem 1 and Example 1 of [8], to prove that X has a continuous version it suffices to show that for each $\phi \in \mathcal{S}(R^d)$ the real Gaussian process $\{\langle X_t, \phi \rangle, t \geq 0\}$ satisfies the Dudley–Fernique condition. Since the covariance functional of X is given by the right-hand side of (2.4) with $n = 1$, for each $\phi \in \mathcal{S}(R^d)$ and $s \leq t$ we have

$$\begin{aligned}
& E[\langle X_t, \phi \rangle - \langle X_s, \phi \rangle]^2 \\
&= \int_{R^d} \phi^2(z) \mu_{ss}(z) dz - \left(\int_{R^d} \phi(z) \mu_{ss}(z) dz \right)^2 \\
&\quad + \int_{R^d} \phi^2(z) \mu_{tt}(z) dz - \left(\int_{R^d} \phi(z) \mu_{tt}(z) dz \right)^2 \\
&\quad - 2 \int_{R^d} \phi(z) \mu_{ss}(z) e^{-\lambda(t-s)} \mathcal{T}_{t-s} \phi(z) dz \\
&\quad + 2 \int_{R^d} \phi(z) \mu_{ss}(z) dz \left[\int_{R^d} \mu_{ss}(z) e^{-\lambda(t-s)} \mathcal{T}_{t-s} \phi(z) dz \right. \\
&\quad \left. + \int_{R^d \times (s, t]} e^{-\lambda(t-r)} \mathcal{T}_{t-r} \phi(x) \mu(dx, dr) \right] \\
&= \int_{R^d} \phi(z) \mu_{ss}(z) [\phi(z) - e^{-\lambda(t-s)} \mathcal{T}_{t-s} \phi(z)] dz \\
&\quad + \int_{R^d} \phi(z) [\phi(z) \mu_{tt}(z) - \mu_{ss}(z) e^{-\lambda(t-s)} \mathcal{T}_{t-s} \phi(z)] dz \\
&\quad + \int_{R^d} \phi(z) \mu_{ss}(z) dz \left\{ \int_{R^d} \mu_{ss}(z) [e^{-\lambda(t-s)} \mathcal{T}_{t-s} \phi(z) - \phi(z)] dz \right\} \\
&\quad + \int_{R^d} \phi(z) \left\{ \int_{R^d} [\mu_{ss}(z) e^{-\lambda(t-s)} (\mathcal{T}_{t-s} \phi(y)) \mu_{ss}(y) \right. \\
&\quad \left. - \mu_{tt}(z) \phi(y) \mu_{tt}(y)] dy \right\} dz \\
&\quad + 2 \int_{R^d} \phi(z) \mu_{ss}(z) dz \int_{R^d \times (s, t]} e^{-\lambda(t-r)} \mathcal{T}_{t-r} \phi(x) \mu(dx, dr).
\end{aligned}$$

The semigroup $\{e^{-\lambda t} \mathcal{T}_t\}$ is differentiable on the Banach space of real functions on R^d having bounded derivatives up to order 2 with norm $\|\phi\|_2 = \sum_{|\alpha| \leq 2} \sup_{x \in R^d} |D^\alpha \phi(x)|$, where $D^\alpha = (\partial/\partial x_1)^{\alpha_1} \cdots (\partial/\partial x_d)^{\alpha_d}$, $|\alpha| = \alpha_1 + \cdots + \alpha_d$. Hence $|\phi(z) - e^{-\lambda(t-s)} \mathcal{T}_{t-s} \phi(z)| \leq M \|\phi\|_2 (t-s)$, where M is a constant. Using this inequality, adding and subtracting $\mu_{ss}(z) \phi(z)$ in the second integral, and $\mu_{ss}(z) \phi(y) \mu_{tt}(y)$ and $\phi(y) \mu_{ss}(y)$ in the fourth integral, we have

$$\begin{aligned}
& E[\langle X_t, \phi \rangle - \langle X_s, \phi \rangle]^2 \\
& \leq M_1 \|\phi\|_2^2 (t-s) + \int_{R^d} \phi^2(z) [\mu_{tt}(z) - \mu_{ss}(z)] dz \\
& \quad + \int_{R^d} \phi(z) \mu_{ss}(z) dz \int_{R^d} \phi(y) [\mu_{ss}(y) - \mu_{tt}(y)] dy \\
& \quad + \int_{R^d} \phi(z) [\mu_{ss}(z) - \mu_{tt}(z)] dz \int_{R^d} \phi(y) \mu_{tt}(y) dy \\
& \quad + 2 \int_{R^d} \phi(z) \mu_{ss}(z) dz \int_{R^d \times (s,t]} e^{-\lambda(t-r)} \mathcal{T}_{t-r} \phi(x) \mu(dx, dr) \\
& \leq M_2 \|\phi\|_2^2 \left\{ (t-s) + \left| \int_{R^d} [\mu_{ss}(z) - \mu_{tt}(z)] dz \right| \right. \\
& \quad \left. + \int_{R^d \times (s,t]} e^{-\lambda(t-r)} \mu(dx, dr) \right\} \\
& \leq M_2 \|\phi\|_2^2 \left\{ (t-s) + 2 \int_{R^d \times (s,t]} e^{-\lambda(t-r)} \mu(dx, dr) \right\} \\
& \leq M_3 \|\phi\|_2^2 (t-s),
\end{aligned}$$

where we have used our assumption on the smoothness of μ and M_1, M_2, M_3 are constants. Hence the Dudley–Fernique condition is satisfied.

By the linearity of X^n and X , proving convergence of the finite-dimensional distributions is equivalent to showing that for all $m \geq 1$, $t_1 \leq \dots \leq t_m$, and $\phi_1, \dots, \phi_m \in \mathcal{S}(R^d)$, $\sum_{i=1}^m \langle X_{t_i}^n, \phi_i \rangle$ converges weakly to $\sum_{i=1}^m \langle X_{t_i}, \phi_i \rangle$ as $n \rightarrow \infty$.

By (2.2) we have

$$\begin{aligned}
\sum_{i=1}^m \langle X_{t_i}^n, \phi_i \rangle &= n^{-1/2} \sum_{i=1}^m \sum_{k=1}^n [\phi_i(B_{t_i}^k) - E\phi_i(B_{t_i}^k)] \\
&= n^{-1/2} \sum_{k=1}^n \sum_{i=1}^m [\phi_i(B_{t_i}^k) - E\phi_i(B_{t_i}^k)].
\end{aligned}$$

Applying the central limit theorem we obtain the convergence of the finite-dimensional distributions.

It remains to show tightness in $D([0, T_0], \mathcal{S}'(R^d))$ and in $D([T_0, \infty), \mathcal{S}'(R^d))$. For this observe that the process X^n can be written as

$$X^n = X^{(1)n} + X^{(2)n},$$

where $X^{(1)n}$ and $X^{(2)n}$ are the fluctuation processes of the empirical distributions of the processes $B^{(1)k}$ and $B^{(2)k}$, $k \geq 1$, respectively, defined by

$$B_t^{(1)k} = \begin{cases} \gamma, & \text{if } t < \tau^k, \\ Z^k + W_{t-\tau^k}^k, & \text{if } t \geq \tau^k, \end{cases}$$

and

$$B_t^{(2)k} = \begin{cases} \gamma, & \text{if } t < \theta^k \\ \tilde{Z}^k + \tilde{W}_{t-\theta^k}^k, & \text{if } t \geq \theta^k, \end{cases}$$

where $\tilde{Z}^k = Z^k + W_{\theta^k - \tau^k}^k$ and $\tilde{W}_t^k = W_{t+(\theta^k - \tau^k)}^k - W_{\theta^k - \tau^k}^k$ for $k \geq 1$.

By Theorem 2.2 of [6], $X^{(1)n}$ converges weakly in $D([0, T_0], \mathcal{S}'(R^d))$ and in $D([T_0, \infty), \mathcal{S}'(R^d))$ and $X^{(2)n}$ converges weakly in $D([0, \infty), \mathcal{S}'(R^d))$; hence $\{(X^{(1)n}, X^{(2)n})\}_{n \geq 1}$ is tight in $D([0, T_0], \mathcal{S}'(R^d)) \times D([0, T_0], \mathcal{S}'(R^d))$ and in $D([T_0, \infty), \mathcal{S}'(R^d)) \times D([T_0, \infty), \mathcal{S}'(R^d))$. Since the limit processes of $X^{(1)n}$ and $X^{(2)n}$ are continuous [6], it then follows (e.g., [1]) that $\{X^{(1)n} + X^{(2)n}\}_{n \geq 1}$ is tight in $D([0, T_0], \mathcal{S}'(R^d))$ and in $D([T_0, \infty), \mathcal{S}'(R^d))$, and the proof is finished.

Observe that X can be considered as an $(\mathcal{S}(R^d) \oplus R)$ -valued process, and its covariance functional is given by the right-hand side of (2.4) with $n = 1$ for $\phi \in \mathcal{S}(R^d) \oplus R$.

For the results on the Markov property and the Langevin equation we consider only $\lambda > 0$, since the case $\lambda = 0$ was discussed in [6].

THEOREM 2.2. (i) *The process $X = \{\langle X_t, \phi \rangle, t \geq 0, \phi \in \mathcal{S}(R^d)\}$ is Markovian if and only if $\mu(R^d \times \{0\}) = 1$.*

(ii) *The process $X^{(a)} \equiv \{\langle X_{a+t}, \phi \rangle, t \in [0, b-a], \phi \in \mathcal{S}(R^d)\}$ is Markovian if and only if $\mu(R^d \times (a, b]) = 0$.*

In both (i) and (ii) the Markov property holds strictly.

(iii) *If $\mu(R^d \times (a, b]) = 0$, the process $X^{(a)}$ satisfies the Langevin equation*

$$dX_t^{(a)} = (\tfrac{1}{2}\Delta - \lambda) X_t^{(a)} dt + dW_t, \quad 0 \leq t \leq b-a, \quad (2.5)$$

where $W = \{W_t, t \geq 0\}$ is an $\mathcal{S}'(R^d)$ -valued Wiener process with covariance functional

$$K_W(s, \phi; t, \psi) = \int_0^{s \wedge t} \langle Q_u \phi, \psi \rangle du, \quad \phi, \psi \in \mathcal{S}(R^d), \quad (2.6)$$

with

$$\langle Q_u \phi, \psi \rangle = \int_{R^d} \mu_{a+u, a+u}(x) [\nabla \phi(x) \cdot \nabla \psi(x) + \lambda \phi(x) \psi(x)] dx \quad (2.7)$$

(\cdot denotes the scalar product in R^d). In particular, if $\mu(R^d \times \{0\}) = 1$ the process X satisfies (2.5)–(2.7) with $a = 0$, $b = \infty$.

Proof. The Markov property holds if and only if given $s < t$, there exists an $\mathcal{S}'(R^d)$ -valued random variable $X_{s,t}$ which satisfies condition (M). Observe that given $r \leq s \leq t$, $\phi, \psi \in \mathcal{S}(R^d)$, from (2.4) we have

$$\begin{aligned} & \text{Cov}[\langle X_r, \phi \rangle, \langle X_t, \psi \rangle] \\ &= \text{Cov}[\langle X_r, \phi \rangle, \langle X_s, e^{-\lambda(t-s)} \mathcal{T}_{t-s} \psi \rangle] \\ &= \int_{R^d} \phi(y) \mu_{rr}(y) dy \int_{R^d \times (s,t]} e^{-\lambda(t-v)} \mathcal{T}_{t-v} \psi(x) \mu(dx, dv). \end{aligned} \quad (2.8)$$

Suppose first that $X_{s,t}$ is of the form $\langle X_{s,t}, \psi \rangle = \langle X_s, e^{-\lambda(t-s)} \mathcal{T}_{t-s} \psi - C_\psi \rangle$, where C_ψ is a constant function which depends on ψ (recall that the process X is well-defined for $\phi \in \mathcal{S}(R^d) \oplus R$). Clearly $\langle X_s, \mathcal{S}(R^d) \oplus R \rangle \subset \langle X_s, \mathcal{S}(R^d) \rangle^\perp$. By (2.8), in order to have the Markov property it suffices to prove

$$\begin{aligned} & \text{Cov}[\langle X_r, \phi \rangle, \langle X_s, C_\psi \rangle] \\ &= \int_{R^d} \phi(y) \mu_{rr}(y) dy \int_{R^d \times (s,t]} e^{-\lambda(t-v)} \mathcal{T}_{t-v} \psi(x) \mu(dx, dv). \end{aligned} \quad (2.9)$$

By (2.4),

$$\begin{aligned} & \text{Cov}[\langle X_r, \phi \rangle, \langle X_s, C_\psi \rangle] \\ &= C_\psi e^{-\lambda(s-r)} \int_{R^d} \phi(z) \mu_{rr}(z) dz \\ &= C_\psi \int_{R^d} \phi(y) \mu_{rr}(y) dy \left[\int_{R^d} \mu_{rs}(y) dy + \int_{R^d \times (r,s]} e^{-\lambda(s-v)} \mu(dx, dv) \right] \\ &= C_\psi (e^{-\lambda(s-r)} - f_{ss}(1)) \int_{R^d} \phi(z) \mu_{rr}(z) dz. \end{aligned} \quad (2.10)$$

From (2.9) and (2.10) we obtain

$$C_\psi = (e^{-\lambda(s-r)} - f_{ss}(1))^{-1} \int_{R^d \times (s,t]} e^{-\lambda(t-v)} \mathcal{T}_{t-v} \psi(x) \mu(dx, dv).$$

For condition (M) to hold C_ψ must not depend on r , and since $\lambda > 0$, this occurs if and only if $\mu(R^d \times (s, t]) = 0$, and since s and t are arbitrary, (M) holds if and only if $\mu(R^d \times \{0\}) = 1$.

Suppose now that $\langle X_{s,t}, \psi \rangle \in \langle X_s, \mathcal{S}(R^d) \rangle^-$ satisfies (M). From our previous calculations we see that provided that we restrict ourselves to $r = s$, then (M) holds for

$$\langle X_{s,t}, \psi \rangle \equiv \langle X_s, \bar{\psi} \rangle \quad (2.11)$$

with

$$\begin{aligned} \bar{\psi} &= e^{-\lambda(t-s)} \mathcal{T}_{t-s} \psi \\ &\quad - (1 - f_{ss}(1))^{-1} \int_{R^d \times (s,t]} e^{-\lambda(t-v)} \mathcal{T}_{t-v} \psi(x) \mu(dx, dv) \quad (\bar{\psi} \in \mathcal{S}(R^d) \oplus R). \end{aligned}$$

We will show that the validity of (M) for $r = s$ implies that (2.11) holds in general.

By assumption there exists a sequence $\{\phi_n\}_n$ in $\mathcal{S}(R^d)$ such that $\langle X_s, \phi_n \rangle$ converges to $\langle X_{s,t}, \psi \rangle$ in L^2 as $n \rightarrow \infty$. Then, using (M) with $r = s$ we have

$$\begin{aligned} E\langle X_s, \bar{\psi} \rangle^2 &= E\langle X_s, \bar{\psi} \rangle \langle X_t, \psi \rangle \\ &= E\langle X_s, \bar{\psi} \rangle \langle X_{s,t}, \psi \rangle \\ &= \lim_{n \rightarrow \infty} E\langle X_s, \bar{\psi} \rangle \langle X_s, \phi_n \rangle \\ &= \lim_{n \rightarrow \infty} E\langle X_t, \psi \rangle \langle X_s, \phi_n \rangle \\ &= \lim_{n \rightarrow \infty} E\langle X_{s,t}, \psi \rangle \langle X_s, \phi_n \rangle \\ &= E\langle X_{s,t}, \psi \rangle^2. \end{aligned}$$

Therefore, again by (M) with $r = s$,

$$\begin{aligned} E|\langle X_{s,t}, \psi \rangle - \langle X_s, \bar{\psi} \rangle|^2 &= E\langle X_{s,t}, \psi \rangle^2 + E\langle X_s, \bar{\psi} \rangle^2 - 2E\langle X_{s,t}, \psi \rangle \langle X_s, \bar{\psi} \rangle \\ &= E\langle X_{s,t}, \psi \rangle^2 + E\langle X_s, \bar{\psi} \rangle^2 - 2E\langle X_s, \bar{\psi} \rangle^2 \\ &= E\langle X_{s,t}, \psi \rangle^2 - E\langle X_s, \bar{\psi} \rangle^2 \\ &= 0, \end{aligned}$$

and hence $\langle X_{s,t}, \psi \rangle = \langle X_s, \bar{\psi} \rangle$ a.s.

But we have seen in the first part of the proof that if

$$\langle X_{s,t}, \psi \rangle = \langle X_s, e^{-\lambda(t-s)} \mathcal{T}_{t-s} \psi + C_\psi \rangle,$$

where C_ψ is a constant, then $\mu(R^d \times \{0\}) = 1$.

To obtain the Langevin equation in (iii) we observe that condition

(SM') holds for $\mathcal{U}_{s,t} = e^{-\lambda(t-s)} \mathcal{T}_{t-s}$. Then we can apply Theorem 2 of [2] (since $\mathcal{S}(R^d) \oplus R$ is a nuclear Fréchet space), which yields the result.

Remarks. (1) In order to compare the present results for $\lambda > 0$ with those for the case $\lambda = 0$ we recall the latter [6]. In this case the fluctuation limit process is always Markovian. If $\mu(R^d \times \{0\}) = 1$, it is strictly Markovian, and it is strictly Markovian on $[a, b]$ if and only if $\mu(R^d \times (a, b]) = 0$. Thus the new behavior exhibited in the case $\lambda > 0$ is that the possibility of death prevents the Markov property. As we have noted in the Introduction, this is a natural result.

(2) The reason for considering convergence on the two spaces $D([0, T_0], \mathcal{S}'(R^d))$ and $D([T_0, \infty), \mathcal{S}'(R^d))$, rather than on $D([0, \infty), \mathcal{S}'(R^d))$, is the lack of a closed martingale on $[0, T_0]$ for the tightness proof [6].

(3) If $\mu(R^d \times (a, b]) = 0$, $X^{(a)}$ is explicitly written as the unique solution of (2.5):

$$\begin{aligned} \langle X_t^{(a)}, \phi \rangle &= e^{-\lambda t} \langle X_0^{(a)}, \mathcal{T}_t \phi \rangle \\ &+ \int_0^t e^{-\lambda(t-s)} \langle dW_s, \mathcal{T}_{t-s} \phi \rangle, \quad 0 \leq t \leq b-a, \phi \in \mathcal{S}(R^d). \end{aligned}$$

3. RELATED DEVELOPMENTS

In the case $\mu(R^d \times \{0\}) = 1$ and $\lambda = 0$, there is an extensive literature on models of the type we have discussed above allowing interactions between the random motions. A small list of recent relevant papers is the following: on fluctuations limits, Dittrich [4], Mitoma [9], Shiga and Tanaka [11], Sznitmann [13], Spohn [12]; on large deviations, Dawson and Gärtner [3], Donsker and Varadhan [5]. It would be interesting to investigate fluctuations and large deviations for such models with space-time creation and/or death ($\mu(R^d \times \{0\}) < 1$, $\lambda > 0$), in particular Markov properties of the fluctuation limits.

ACKNOWLEDGMENT

I am grateful to L. G. Gorostiza for suggesting this problem and for his guidance during the course of this research.

REFERENCES

1. D. J. ALDOUS, Weak convergence and the general theory of processes, unpublished manuscript, 1981.

2. T. BOJDECKI AND L. G. GOROSTIZA, Inhomogeneous infinite dimensional Langevin equations, *Stochastic Anal. Appl.* **6** (1988), 1–9.
3. D. A. DAWSON AND J. GÄRTNER, Large deviations from the McKean–Vlasov limit for weakly interacting diffusions, *Stochastics* **20** (1987), 247–308.
4. P. DITTRICH, A stochastic particle system: Fluctuations around a nonlinear reaction-diffusion equation, *Stochastic Process. Appl.* **30** (1988), 149–164.
5. M. D. DONSKER AND S. R. S. VARADHAN, Large deviations from a hydrodynamic scaling limit, *Comm. Pure Appl. Math.* **42** (1989), 243–270.
6. B. FERNÁNDEZ, Markov properties of the fluctuation limit of a particle system, *J. Math. Anal. Appl.* **149** (1990), 160–179.
7. K. ITO, Distribution-valued processes arising from independent Brownian motions, *Math. Z.* **182** (1983), 17–33.
8. I. MITOMA, On the norm continuity of \mathcal{S}' -valued Gaussian processes, *Nagoya Math. J.* **82** (1981), 209–220.
9. I. MITOMA, Tightness of probabilities on $C([0, 1], \mathcal{S}')$ and $D([0, 1], \mathcal{S}')$, *Ann. Probab.* **11** (1983), 989–999.
10. I. MITOMA, Generalized Ornstein–Uhlenbeck processes having a characteristic operator with polynomial coefficients, *Probab. Theory Relat. Fields* **76** (1987), 533–555.
11. T. SHIGA AND H. TANAKA, Central limit theorem for a system of Markovian particles with mean field interactions, *Z. Wahrsch. Verw. Gebiete* **69** (1985), 439–459.
12. H. SPOHN, Equilibrium fluctuations for interacting Brownian particles, *Comm. Math. Phys.* **103** (1986), 1–33.
13. A. S. SZNITMANN, Nonlinear reflecting diffusion process and the propagation of chaos and fluctuations associated, *J. Funct. Anal.* **56** (1984), 311–336.